

TYPE: U.S.A. California: Fresno Co., E side of summit of Parkfield Grade, *Hall 11769* (holotype, UC).

*Acknowledgments.* Supported in part by NSF grant BSR 89-08963. I thank John Strother, Ron Hartman, John Semple, Greg Brown, J. P. Saroyan, and D. R. Parnell for assistance.

#### Literature Cited

- Canby, M. L. 1927. The genus *Corethrogyne* in southern California. *Bull. S. Calif. Acad. Sci.* 26: 8-16.
- Gottlieb, L. D. 1984. Genetics and morphological evolution in plants. *Amer. Naturalist* 123: 681-709.
- Howell, J. T. 1929. A systematic study of the genus *Lessingia* Cham. *Univ. Calif. Publ. Bot.* 16: 1-44.
- Jackson, R. C. & C. T. Dimas. 1981. Experimental evidence for systematic placement of the *Happypappus phyllocephalus* complex (Compositae). *Syst. Bot.* 6: 8-14.
- Keck, D. D. 1956. *Benitoa*, a new genus of Compositae from California. *Leaf. W. Bot.* 8: 25-40.
- . 1959. *Corethrogyne*. Pp. 1204-1207 in P. A. Munz, *A California Flora*. Univ. California Press, Berkeley & Los Angeles.
- Spence, W. L. 1963. A biosystematic study of the genus *Lessingia* Cham. (Compositae). Unpublished Ph.D. dissertation, University of California, Berkeley. (University Microfilms order no. 64-2136.)